

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

Vol. 149, No. 3, 1987  
December 31, 1987

BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS  
Pages 960-968

EXPRESSION OF CHIMERIC RECEPTOR COMPOSED OF IMMUNOGLOBULIN-DERIVED  
V REGIONS AND T-CELL RECEPTOR-DERIVED C REGIONS

Yoshihisa Kawanishi<sup>1</sup>, Yoshihiro Asakura<sup>1</sup>, Naoko Utsunomiya<sup>2</sup>,  
Mamoru Nakanishi<sup>2</sup>, Yohji Arata<sup>3</sup>, Seiga Itoh<sup>3</sup>,  
Fumihiko Nagase<sup>4</sup> and Yoshikazu Kurosawa<sup>1\*</sup>

<sup>1</sup>Institute for Comprehensive Medical Science, Fujita-Gakuen Health  
University, Toyouke, Aichi, 470-11

<sup>2</sup>Faculty of Pharmaceutical Science, University of Tokyo, Hongo,  
Bunkyo-ku, Tokyo, 113

<sup>3</sup>Tokyo Research Laboratories, Kyowa Hakko Co., Asahimachi,  
Machida, Tokyo, 194

<sup>4</sup>Department of Immunology, Nagoya University School of Medicine,  
Tsurumai, Showa-ku, Nagoya, 466, Japan

Received November 12, 1987

**SUMMARY:** Chimeric genes composed of immunoglobulin (Ig)-derived variable (V) regions and T-cell receptor (TCR)-derived constant (C) regions were constructed. The V<sub>L</sub> and V<sub>H</sub> genes showing anti-phosphorylcholine (PC) activity were used in this study. Two pairs of chimeric genes, V<sub>L</sub>-C<sub>β</sub> and V<sub>H</sub>-C<sub>α</sub> genes, and V<sub>L</sub>-C<sub>α</sub> and V<sub>H</sub>-C<sub>β</sub> genes, were inserted into an expression vector containing both Ecogpt and neo genes, and transfected into EL4 cells. Cells which express both chimeric receptor molecules were established. The activity of the transformants to the antigen was examined by using stopped-flow fluorometry. An increase in the concentration of cytoplasmic calcium ion was observed after addition of *Staphylococcus pneumoniae* R36A bacteria grown in the choline-containing medium which express PC molecules, but not after the PC-negative bacteria grown in the ethanolamine-containing medium. © 1987 Academic Press, Inc.

TCR is a heterodimer of disulfide-linked α and β chains (1). The essential features of both chains are rather similar to those of Ig(2). N-terminal regions of both chains are variable and form a dimeric antigen-combining site (3). In contrast to the recognition of antigen by antibody, where the antibody can bind to free antigen, the TCR does not recognize antigen alone but only in association with MHC molecules (4). To determine whether the difference in antigen recognition between T and B

\* To whom correspondence should be addressed.

**Abbreviations:** Ig, immunoglobulin; TCR, T-cell receptor; V, Variable; C, constant; H, heavy; L, light; PC, phosphorylcholine; ECO-MOL, ecotropic Moloney virus; LTR, long terminal repeat.

0006-291X/87 \$1.50

Copyright © 1987 by Academic Press, Inc.

All rights of reproduction in any form reserved.

960

Vol. 149, No. 3,

cells derived  
constructed  
C regions.  
study. The  
myelomas are  
germline V<sub>H</sub>  
the anti-PC  
were established  
accompanied  
The activity  
stopped-flow  
on the trans  
T cell activ

Active  
containing  
myeloma S10  
The C gene  
C57BL/6 mouse  
M. Davis (St.  
Moloney virus  
M. Ishimoto  
previously  
plasmid show  
resistance to  
sequence of  
method (13).  
Transfection  
Total RNA  
Poly A-conta  
(15). Nor  
antibody (17  
(Israel) and  
was carried  
cells/ml) we  
25°C. Stop  
flow spectro  
system (8).  
filter (whic  
to come into

Figure  
genes: V<sub>L</sub>-C<sub>β</sub>  
T cells, we  
translation  
(20). we de

cells derives from the structural difference in V regions of TCR and Ig, we constructed chimeric genes composed of Ig-derived V regions and TCR-derived C regions. The  $V_L$  and  $V_H$  gene showing anti-PC activity were used in this study. The V genes of the light (L) and heavy (H) chains of anti-PC myelomas and hybridomas have been well characterized (5,6). Only one germline  $V_H$  gene (5) and three germline  $V_L$  genes (6) are responsible for the anti-PC specificity. Cells which express chimeric receptor molecules were established. It is known that the activation of T lymphocytes is accompanied by an increase in the concentration of cytosolic calcium (7). The activity of the transformants to the antigen was examined by using stopped-flow fluorometry (8). The chimeric receptor molecules expressed on the transformants had the capacity to react with PC antigen and trigger T cell activation.

#### MATERIALS AND METHODS

Active  $V_H$  gene was isolated from myeloma TEPC15 by using  $J_H$  gene-containing DNA fragment as a probe (5). Active  $V_L$  gene isolated from myeloma S107 was donated by M.D. Scharif (Albert Einstein College) (9). The C genes of T-cell receptor  $\alpha$  and  $\beta$  chains were isolated from C3H and C57BL/6 mouse DNA by using cDNAs of  $\alpha$  and  $\beta$  chains (2,10) donated by M. Davis (Stanford University) as probes, respectively. The ecotropic Moloney virus (ECO-MOL) long terminal repeat (LTR) fragment was donated by M. Ishimoto (Kyoto University) (11). Plasmid pAGE145 was described previously (12). The bacteria and the eukaryotic cells containing this plasmid show kanamycin resistance at a concentration of 10  $\mu$ M and G418 resistance at a concentration of 1 mg/ml, respectively. The nucleotide sequence of the constructed plasmids was determined by the M13-dideoxy method (13). EL4 cell was obtained from T. Taniguchi (Osaka University). Transfection experiments were done by protoplast fusion as described (14). Total RNA was extracted from cells by the Guanidinium-CsCl method (15). Poly A-containing RNA was purified with the oligo (dT) cellulose column (15). Northern hybridization was performed as described (16). Anti- $V_H$  antibody (17) and anti-T15 idiotype antibody (18) were donated by D. Civol (Israel) and C. Heusser (Basel), respectively. Stopped-flow fluorometry was carried out as follows. Fura 2-loaded T-cells (19) (final  $10^6$  cells/ml) were mixed with the target bacteria (final  $4 \times 10^8$  cells/ml) at 15°C. Stopped-flow fluorescence was measured with a Union Giken stopped-flow spectro-photometer RA401 in combination with a microcomputer RA-450 system (8). Excitation wavelength: 335nm. We used a Hoya Y46 cut filter (which allows the emitted light with wavelength longer than 460 nm to come into the detector) in the fluorescence measurements.


#### RESULTS

Figure 1 shows the construction procedures for four kinds of chimeric genes:  $V_L$ -C $\beta$ ,  $V_H$ -C $\alpha$ ,  $V_L$ -C $\alpha$  and  $V_H$ -C $\beta$ . To express the chimeric genes in T cells, we used the ECO-MOL LTR as a transcriptional promoter. Since translation begins at the 5'-proximal AUG triplet in eukaryotic mRNA (20), we determined the nucleotide sequence between the promoter and

Vol. 149, No. 3, 1987

BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS

Vol. 149, N. 3, 1:



**Fig. 1.** Construction of the chimeric genes. (a) Restriction maps of the clones. The DNA fragments used for the construction of plasmids are indicated by thick lines. Restriction Enzymes: E, EcoRI; B, BamHI; H, HindIII; X, XbaI; S, SmaI; Bg, BglII; P, PstI; K, KpnI; C, ClaI; Sa, SacI; S, SmaI. (b) For the construction of pSVG-MOLV\_Cg37, the ClaI-SmaI fragment of the ECO-MOLV LTR(LTR-I) was connected to the BamHI-BamHI fragment of T15V (V<sub>II</sub>-1), after the BamHI site at the 5' upstream of V<sub>II</sub> gene was changed to a flush end with Klenow enzyme. The resulting ClaI-BamHI fragment was connected to the BamHI-BamHI fragment of TCR α chain as in the same polarity, and inserted into plasmid pSV2gpt. The EcoRI-ClaI region at the upstream of the LTR sequence was derived from pBR322 DNA. ; For the construction of pSVG-MOLV\_Cg12, the LTR-I fragment was connected to the SmaI-BglII fragment of S107V (V<sub>I</sub>). The resulting fragment was connected to the BamHI fragment of C<sub>0</sub> gene after changing the BglII site into BamHI site by using a BamHI linker, and inserted into plasmid pSV2gpt. ; For the construction of pSVG-MOLV\_Cg11, the ClaI-Kpn fragment of the ECO-MOLV

the leader of ATG sequence for transfection cell (21), was MOLV\_C<sub>0</sub>(neo), protoplast of medium (23). gpt and neo pV<sub>H</sub>C<sub>0</sub>V<sub>II</sub>C<sub>0</sub><sup>3</sup> chimeric gene in the gpt<sup>+</sup> transfected respectively. transformants Northern hybridization cells expressed EL4-5 contained neo resistant 1.45 kb with a

LTR(LTR-II) after changing enzyme, which fragment was changing respectively the construction pSVG-MOLV\_C<sub>0</sub>. Instead of whose structure resistance pSV2gpt. MOLV\_C<sub>0</sub> (neo) construction DNA, pAGE14: polylinker designation: splicing site thymidine kinase of P1 and MOLV\_C<sub>0</sub>Al1 site of pAGE14: pSVG-MOLV\_C<sub>0</sub> resulting fragment BglII fragment pSVG-MOLV\_C<sub>0</sub> pAGE-V<sub>I</sub>C<sub>0</sub>18. ClaI-digestion pV<sub>I</sub>C<sub>0</sub>V<sub>II</sub>C<sub>0</sub>16. pV<sub>I</sub>C<sub>0</sub>V<sub>II</sub>C<sub>0</sub>14. The overall hybridization sizes of each

the leader of  $V_H$  and  $V_L$  genes in the constructed plasmids. There were no ATG sequences in the 5' upstream regions (Fig.2). In an initial trial for transformation experiments, the EL4 cells, presumed to be a helper T cell (21), were transfected with two plasmid DNAs: pSVC-MOLV $C_{\alpha}37$  and pSVC-MOLV $C_{\beta}$ (neo), or pSVG-MOLV $C_{\alpha}12$  and pSVG-MOLV $C_{\beta}$ (neo), simultaneously, by protoplast fusion (14) and selected in the gpt (22) and neo selection medium (23). However, we did not obtain any transformants showing both gpt and neo resistance. We constructed the plasmids: pV $C_{\beta}V_L C_{\alpha}1$ , pV $C_{\beta}V_L C_{\alpha}3$ , pV $C_{\beta}V_H C_{\alpha}14$  and pV $C_{\beta}V_H C_{\alpha}16$ , containing two chimeric genes in each plasmid DNA (Fig.1c). From  $4 \times 10^6$  EL4 cells, in the gpt selection medium, we obtained 7, 22, 13 and 6 transformants transfected with pV $C_{\beta}V_L C_{\alpha}1$ , pV $C_{\beta}V_L C_{\alpha}3$ , pV $C_{\beta}V_H C_{\alpha}14$  and pV $C_{\beta}V_H C_{\alpha}16$ , respectively. In the neo selection medium, we obtained 2, 10, 9 and 3 transformants transfected with the above plasmid DNAs, respectively. By Northern hybridization (16) with the  $V_H$  and  $V_L$  probes, we searched for cells expressing both chimeric genes, and obtained two transformants: EL4-5 containing pV $C_{\beta}V_L C_{\alpha}3$  and EL4-14 containing pV $C_{\beta}V_H C_{\alpha}14$ , as neo resistant cells. EL4-5 gave a band at 1.35 kb with the  $V_H$  probe and at 1.45 kb with the  $V_L$  probe (Fig.3). EL4-14 gave a band at 1.45 kb with

LTR(LTR-II) was connected to the BamHI-HindIII fragment of Tbv ( $V_H$ -11) after changing the KpnI and the BamHI sites into flush ends with Klenow enzyme, which creates a BamHI site at the junction point. The resulting fragment was connected to the KpnI-HindIII fragment of  $C_{\beta}$  gene after changing the KpnI and HindIII sites into BamHI and BamHI sites, respectively, by using linkers, and inserted into plasmid pSV2gpt. For the construction of pSVG-MOLV $C_{\beta}A1$ , the protocol was the same as that of pSVC-MOLV $C_{\alpha}12$  except for the change of the BglII site into HindIII site instead of BamHI site. Independently, we constructed four other plasmids whose structures are essentially the same as shown in (b) except for neo resistance gene in place of EcoRpt gene by using plasmid pSV2neo instead of pSV2gpt. We named them pSVG-MOLV $C_{\alpha}$ (neo), pSVG-MOLV $C_{\beta}$ (neo), pSVG-MOLV $C_{\alpha}$ (neo), pSVG-MOLV $C_{\beta}$ (neo), respectively. (c) For the construction of the plasmids containing two chimeric genes in one plasmid DNA, PAGE145(31) was used as a vector. The LTR in the ECO-MOLTR and a polylinker shown in Fig.2 was inserted into the SmaI site. The designations are as follows: SpBC, ARG, ASE and Ark are  $\beta$ -globin splicing signal, A-globin polyA signal, SV40 early gene polyA signal and thymidine kinase polyA signal, respectively. Pl and Prk are the promoters of Pl and thymidine kinase genes. The BamHI-BamHI fragment of pSVC-MOLV $C_{\beta}A1$  containing the chimeric  $V_H$ - $C_{\beta}$  gene was inserted into the BamHI site of PAGE145, resulting in the plasmid PAGE-V $C_{\beta}6$ . The ClaI-digested pSVC-MOLV $C_{\alpha}12$  was connected to the ClaI-digested PAGE-V $C_{\beta}6$ , resulting in two clones: pV $C_{\beta}V_L C_{\alpha}1$  and pV $C_{\beta}V_L C_{\alpha}3$ . The SacI-BglII fragment of PAGE145 was replaced by the SacI-BamHI fragment of pSVC-MOLV $C_{\beta}A1$  containing the chimeric  $V_L$ - $C_{\beta}$  gene, resulting in plasmid PAGE-V $C_{\beta}18$ . The ClaI-digested pSVC-MOLV $C_{\alpha}37$  was connected to the ClaI-digested PAGE-V $C_{\beta}18$ , resulting in two plasmids: pV $C_{\beta}V_H C_{\alpha}14$  and pV $C_{\beta}V_H C_{\alpha}16$ . The sizes of the four plasmids: pV $C_{\beta}V_L C_{\alpha}1$ , pV $C_{\beta}V_L C_{\alpha}3$ , pV $C_{\beta}V_H C_{\alpha}14$ , and pV $C_{\beta}V_H C_{\alpha}16$  are 23, 23, 21.6 and 21.6 kb, respectively. The overall structures are confirmed by restriction mapping. Southern hybridization and nucleotide sequencing of the connected regions. The sizes of each plasmid and gene shown in (b)(c) and (d) are arbitrary.

Vol. 149, No. 3, 1987

BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS

Vol. 149, No. 3, 198




Fig.2. Nucleotide sequence from the promoter region to the first intron of the constructed V<sub>H</sub> and V<sub>L</sub> genes. The nucleotide sequence of the promoter was as described(11). The junction points of the LTR and V genes are at the SmaI or KpnI site in the ECO-MOI LTR, the BamHI site in the 5' upstream of I15 V<sub>H</sub> gene and the StuI site in the 5' upstream of S107 V<sub>L</sub> gene as described in the legend of Fig.1. Since a polylinker was inserted into the SmaI site in plasmid PAGE145 (12), there are 25 extra nucleotides at the boundary of the LTR and 5' V<sub>H</sub> in pMOI-V<sub>H</sub> 6. The determined sequences are the same as the published sequence(5) except for a two nucleotide deletion in the 5' untranslated region of the V<sub>H</sub> gene. The position of the initiation codon in S107 V<sub>L</sub> gene is estimated from the locations of the octamer ATTTGCAT(35) and TATA-like sequences whose portions were removed from the constructed plasmids.

the V<sub>H</sub> probe and at 1.25 kb with the V<sub>L</sub> probe. The size of each band indicates that RNA splicing occurred between an Ig-derived J gene and a TCR-derived C gene. The size differences should reflect those of the 3' end structure of the constructed genes(Fig.1).




Fig.3. Northern hybridization of the transformants with the V<sub>H</sub> and V<sub>L</sub> probe. PolyA-containing RNA was prepared from each transformant. Two micrograms of RNA were subjected to 1% agarose gel electrophoresis. The size was estimated using 28S and 18S rRNA as size markers. Hybridization was carried out with (a)(c) the V<sub>H</sub> probe (which corresponds to V<sub>H</sub>II in Fig.1a), and (b)(d) the V<sub>L</sub> probe (which corresponds to V<sub>L</sub> in Fig.1a). (a)(d)EL4-5, (c)(b)EL4-14.

To determine chimeric receptors used two monoclonal idiotype antisera reacted with crossreactivity (24), or some antibody were problems, we antigen without

It is known increase in the target cells, then, calcium calcium influx. This increase is using stopped-source of PC in bacteria (25) molecules,<sup>9</sup> and bacteria grown in both EL4-5 and the T cells with and that EL4-5 bacteria. The presumably expressed PC antigen and

The TCR is Lyt2/T8 and L3T4 interactions between groups (30,31) TCR alone may Transfection allow the reciprocal allelic MHC presentation difference in the structural computer analyses

## IMUNICATIONS

Vol. 149, No. 3, 1987

## BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS

CTCACTCGG

AGCACTCAG

C GT(intron)

Met  
AGCAGG ATG

intron  
of the  
genes  
the 5'  
17 V  
inserted  
otides  
imined  
a two  
The  
in the  
whose

each band  
ne and a  
of the 3'

nd V<sub>H</sub>  
Two  
The  
zation  
in in  
g.la).

To determine whether translation of mRNAs and expression of the chimeric receptor molecules on the cell surface occurred in these cells, we used two monoclonal antibodies: anti-V<sub>H</sub> antibody (17) and anti-T15 idiotype antibody (18). However, the EL4 used as a recipient cell itself reacted with anti-T15 idiotype antibody, presumably due to the crossreactivity to Thy-1 which shares the same epitope as T15 idiotype (24), or some other surface molecules, and the results with anti-V<sub>H</sub> antibody were ambiguous because of the high background. Because of these problems, we directly examined the activity of the transformants to the antigen without identification of the proteins.

It is known that the activation of T lymphocytes is accompanied by an increase in the concentration of cytosolic calcium (7). After binding to target cells, T lymphocytes first increase their membrane fluidity and, then, calcium is released from intracellular stores (8). After that, calcium influx occurs from the external medium into T lymphocytes (8). This increase in the concentration of cytoplasmic Ca<sup>2+</sup> ion can be traced by using stopped-flow fluorometry (8) with a fluorescent probe. As the source of PC antigen, we used heat-killed *Staphylococcus pneumoniae* R36A bacteria (25) grown in the chorine-containing medium which express PC molecules,<sup>9</sup> and as the negative control, we used heat-killed PC-negative bacteria grown in arbanolamine-containing medium (27). Figure 4 shows that in both EL4-5 and EL4-14 cells, calcium influx was observed after mixing the T cells with PC-positive bacteria but not with PC-negative bacteria, and that EL4 itself did not react with either PC-positive or PC-negative bacteria. These results indicate that the chimeric receptor molecules presumably expressed in the transformants have the capacity to react with PC antigen and trigger T cell activation.

## DISCUSSION

The TCR is associated with the T3 complex(28). Furthermore, the L2/2/T8 and L24/T4 surface glycoproteins are thought to be involved in the interactions between T cells and their target cells (29). However, two groups (30,31) have presented direct evidence that the  $\alpha$  and  $\beta$  chains of TCR alone may define the dual, namely antigen and MHC, specificity. Transfection and expression of the genes for a particular  $\alpha$  and  $\beta$  pair allow the recipient cell to respond to the combination of antigen and allelic MHC product recognized by the donor T cell (30,31). The difference in antigen recognition between B and T cells may originate in the structural difference in Ig V regions and TCR V regions, although computer analyses (32), based on published sequence data, indicate that the




Fig. 4. Stopped-flow fluorometry traces indicating the effects of the antigens on the T-cells with chimeric receptors.

The data are shown as the average of 15 times measurements.

- (a) Effect of  $PC^+$  on the fluorescence of fura 2-loaded EL4-5.
- (b) Effect of  $PC^-$  on the fluorescence of fura 2-loaded EL4-5.
- (c) Effect of  $PC^+$  on the fluorescence of fura 2-loaded EL4-14.
- (d) Effect of  $PC^-$  on the fluorescence of fura 2-loaded EL4-14.
- (e) Effect of  $PC^+$  on the fluorescence of fura 2-loaded EL4.
- (f) Effect of  $PC^-$  on the fluorescence of fura 2-loaded EL4.

$\alpha$  and  $\beta$  chains of TCR are organized into immunoglobulin-like domains consisting of multistranded antiparallel  $\beta$ -sheet bilayers. Our present experiments indicate that the chimeric receptor composed of Ig-derived V regions and TCR-derived C regions has the capability to trigger T cell

activation with influx is an helper and cytotoxic the ability of cells. In any antigens described in the Recently, chimeric genes Although a chimeric to form an appropriate secreted, it reason for for and  $V_L$ - $C_H$ , a Ig's character without C domain

We thank C. Heusser and Dr. R. K. We thank Dr. R. K. are grateful for their encouragement for the Ministry of Health University of the Uehara Science

1. Kappler, J., 727-737.
2. Hedrick, S.M. Nature 308, 4, 529-591.
3. Kronenberg, 4, 529-591.
4. Schwartz, R.
5. Crews, S., 59-66.
6. Gearhart, P., 29-34.
7. Oertgen, H.C. 40, 583-590.
8. Utsunomiya, Proc. Natl. Acad. Sci. USA 85, 100-104.
9. Kwan, S.-P., J. Exp. Med. 171, 100-104.
10. Chien, Y., Davis, M.M.
11. Adachi, A., & Ishimoto, J.



## COMMUNICATIONS

Vol. 149, No. 3, 1987

## BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS

activation with the antigen alone, without MHC molecules. Since calcium influx is an early transmembrane event (8), we are now examining whether helper and cytolytic functions occur with antigens. If this is the case, the ability of Ig to bind to free antigens could be transferred into T cells. In future, it might become possible for T cells recognizing any antigens without MHC restriction to be produced by the technique described in this paper.

Recently, Cascoigne et al. (33) reported the expression of reverse chimeric genes composed of TCR-derived V regions and Ig-derived C regions. Although a chimeric protein  $V_{\alpha}-C_{\gamma 2b}$  combined with a normal  $\lambda$  light chain to form an apparently normal tetrameric immunoglobulin molecule that was secreted, it did not combine with a chimeric  $V_{\beta}-C_{\kappa}$  chain (33). The reason for formation of our chimeric chains:  $V_{H}-C_{\beta}$  and  $V_{L}-C_{\alpha}$ , and  $V_{H}-C_{\alpha}$  and  $V_{L}-C_{\beta}$ , although not directly demonstrated, might be related to the Ig's character in that  $V_{H}$  and  $V_{L}$  domains can be intrinsically associated without C domains to form an  $F_{\gamma}$  fragment (34).

## ACKNOWLEDGEMENTS

We thank Drs. M. Davis, A. Ishimoto, H. Sero, M. D. Scharff, D. Civol, C. Heusser and M. Potter for providing us the materials used in this study. We thank Dr. R. Kubo for helpful discussions to prepare this manuscript. We are grateful to Drs. M. Hatakeyama and T. Taniguchi for teaching us the transfection method. We also thank Drs. Y. Takagi, I. Ishiguro and K. Fujita for their encouragement. This work was supported in part by grants from the Ministry of Education, Science and Culture of Japan; Fujita-Gakuen Health University; Japan Private School Promotion Foundation (to Y.K.) and the Uehara Science Foundation (to Y.K.).

## REFERENCES

1. Kappler, J., Kubo, R., Haskins, K., White, J. & Marrack, P. (1983) *Cell* **34**, 727-737.
2. Hedrick, S.M., Nielsen, E.A., Kavaler, J., Cohen, D.L. & Davis, M.M. (1984) *Nature* **308**, 153-158.
3. Kronenberg, M., Siu, G., Hood, L.E. & Shastri, N. (1986) *Ann. Rev. Immunol.* **4**, 529-591.
4. Schwartz, R.H. (1985) *Ann. Rev. Immunol.* **3**, 237-261.
5. Crews, S., Griffin, J., Huang, H., Calame, K. & Hood, L. (1981) *Cell* **25**, 59-66.
6. Gearhart, P.J., Johnson, N.D., Douglas, R. & Hood, L. (1981) *Nature* **291**, 29-34.
7. Oettingen, H.C., Terhorst, C., Cantley, L.C., & Rosoff, P.M. (1985) *Cell* **40**, 583-590.
8. Utsunomiya, N., Tsuboi, M., & Nakanishi, M. (1986) *Proc. Natl. Acad. Sci. USA* **83**, 1877-1880.
9. Kwan, S.-P., Rudikoff, S., Seidman, J.G., Leder, P. & Scharff, M.D. (1981) *J. Exp. Med.* **154**, 1366-1370.
10. Chien, Y., Becker, D.M., Lindsten, T., Okamura, M., Cohen, D.L. & Davis, M.M. (1984) *Nature* **312**, 31-35.
11. Adachi, A., Sakai, K., Kitamura, N., Nakanishi, S., Niwa, O., Matsuyama, M. & Ishimoto, A. (1984) *J. Virol.* **50**, 813-821.

Vol. 149, No. 3, 1987

BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS

12. Mizukami, T. & Ito, S. (1987) *J. Biochem.* **101**, 1307-1310.
13. Messing, J., Crea, R. & Seeburg, P.H. (1981) *Nucl. Acid. Res.* **9**, 309-321.
14. Hatakeyama, M., Minamoto, S., Uchiyama, T., Hardy, R.R., Yamada, G. & Taniguchi, T. (1985) *Nature* **318**, 467-471.
15. Maniatis, T., Fritsch, E.F. & Sambrook, J. eds (1982) *Molecular cloning, a laboratory manual*, Cold Spring Harbor laboratory.
16. Thomas, P.S. (1980) *Proc. Natl. Acad. Sci. USA* **77**, 5201-5205.
17. Eshhar, Z., Gligi, O., Givol, D. & Ben-Neriah, Y. (1983) *Eur. J. Immunol.* **13**, 533-540.
18. Carny, J., Heusser, C., Wallich, R., Hammerling, G.J. & Eardley, D.D. (1982) *J. Exp. Med.* **156**, 719-730.
19. Crynkiewicz, G., Poenie, M. & Tsien, R.Y. (1985) *J. Biol. Chem.* **260**, 3440-3450.
20. Kozak, M. (1986) *Cell* **44**, 283-292.
21. Farrar, J.J., Benjamin, W.R., Hilfiker, M.L., Howard, M., Farrar, W.L. & Fuller-Farrar, J. (1982) *Immunol. Rev.* **63**, 129-166.
22. Mulligan, R.C. & Berg, P. (1980) *Science* **209**, 1422-1427.
23. Southern, P.J. & Berg, P. (1982) *J. Mol. Appl. Genet.* **1**, 327-341.
24. Pillemer, E. & Weissman, I.L. (1981) *J. Exp. Med.* **153**, 1068-1074.
25. Chang, S.P., Perlmutter, R.M., Brown, M., Heusser, C.H., Hood, L. & Rittenberg, M.R. (1984) *J. Immunol.* **132**, 1550-1555.
26. Mosser, J.L. & Tomasz, A. (1970) *J. Biol. Chem.* **245**, 287-298.
27. Tomasz, A. (1968) *Proc. Natl. Acad. Sci. USA* **59**, 86-93.
28. Reinherz, E.L., Meuer, S.C., Fitzgerald, K.A., Hussey, R.E., Hodgdon, J.C., Acuto, O. & Schlossman, S. (1983) *Proc. Natl. Acad. Sci. USA* **80**, 4104-4108.
29. Biddison, W.E., Rao, P.E., Talle, M.A., Goldstein, G. & Shaw, S. (1981) *J. Exp. Med.* **159**, 793-797.
30. Denhio, Z., Haas, W., Weiss, S., McCubrey, J., Kiefer, H., von Boehmer, H. & Steinmetz, M. (1986) *Nature* **320**, 232-238.
31. Saito, T., Weiss, A., Miller, J., Norcross, M.A., & Germain, R.N. (1987) *Nature* **325**, 125-130.
32. Novotny, J., Tonegawa, S., Saito, H., Krantz, D.M., & Eisen, H.N. (1986) *Proc. Natl. Acad. Sci. USA* **83**, 142-146.
33. Gascoigne, N.R.J., Goodnow, C.C., Dudzik, K.L., Oi, V.T. & Davis, M.M. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 2936-2940.
34. Novotny, J., & Haber, E. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 4592-4596.
35. Parslow, T.G., Blair, D.L., Murphy, W.J. & Granner, D.K. (1984) *Proc. Natl. Acad. Sci. USA* **81**, 2650-2655.

Vol. 149, No. 3, 1987  
December 31, 1987THE EARLY COM  
IN 2T.J. Koerner, <sup>2</sup> TLaboratory  
Pathology; <sup>1</sup> Deper  
Duk

Received November

**SUMMARY.** Tri-  
lipopolysaccharide  
mRNA for the ear  
multiple pathways  
mRNA for JE was  
was not (Introna et  
regulation of the  
present evidence th  
the JE gene. By  
induced transcripti  
gene in LPS-treat  
gene is regulated  
mechanistic pathwa  
upon the cell type:  
© 1987 Academic Press.

**INTRODUCTION.**  
defense mechanism  
potent changes in

1. Supported in  
grant #1783 f  
RJR Nabisc .
2. To whom cor  
Biology of Lei
3. Present Addr  
Cleveland, OH
4. Present Addre  
Abbreviations  
platelet-deriv